MATRIX ALGEBRA MAGNUS JANSSON DEADLINE: 2016–05–25, 10.00

Homework # 8

1. Let

$$A = \begin{bmatrix} 1 & 0 \\ 0 & 2 \end{bmatrix}$$
$$B = \begin{bmatrix} 0 & 2 \\ 0 & 0 \end{bmatrix}$$

Determine F(A), F(B), F(A+B), and show that F(A+B) is a proper subset of F(A) + F(B).

- 2. Show that $A \in M_2(R)$ is positive stable if and only if tr(A) > 0 and det(A) > 0.
- 3. Let $B \in M_n$ be a matrix none of whose eigenvalues μ_i is equal to 1, and define

 $A = (B + I)(B - I)^{-1}$

Show that the eigenvalues of A are given by

$$\lambda_i = \frac{\mu_i + 1}{\mu_i - 1}$$

Prove that $\operatorname{Re}(\lambda_i) < 0$ if and only if $|\mu_i| < 1$. Conclude that A is negative stable if and only if B is a convergent matrix.

4. Use the transformation in the previous exercise to derive the discrete time version of the Lyapunov equation from the continuous time counterpart. (That is, derive $A^*GA - G = -H$ from $GA + A^*G = -H$, where the last negative sign is because we discuss negative stability here.)