


# Think about this:

$$\mathbf{F} = q\mathbf{v} \times \mathbf{B}$$



What happens if you add an electric field  $\mathbf{E}$ ?



# Last lecture (3)

- Solar activity
- Solar wind – basic facts

# Today's lecture (4)

- Solar wind – magnetic structure
- Ionosphere
  - layers
  - radio wave reflection
  - electrical conductivity in magnetized plasma



# Today

<u>Activity</u>	<u>Date</u>	<u>Time</u>	<u>Room</u>	<u>Subject</u>	<u>Litterature</u>
L1	2/9	10-12	Q33	Course description, Introduction, The Sun 1, Plasma physics 1	<b>CGF</b> Ch 1, 5, (p 110-113)
L2	3/9	15-17	Q31	The Sun 2, Plasma physics 2	<b>CGF</b> Ch 5 (p 114-121), 6.3
L3	9/9	10-12	Q33	Solar wind, The ionosphere and atmosphere 1, Plasma physics 3	<b>CGF</b> Ch 6.1, 2.1-2.6, 3.1-3.2, 3.5, <b>LL</b> Ch III, Extra material
T1	11/9	10-12	Q34	Mini-group work 1	
L4	16/9	15-17	Q33	The ionosphere 2, Plasma physics 4	<b>CGF</b> Ch 3.4, 3.7, 3.8
L5	18/9	15-17	Q21	The Earth's magnetosphere 1, Plasma physics 5	<b>CGF</b> 4.1-4.3, <b>LL</b> Ch I, II, IV.A
T2	23/9	10-12	Q34	Mini-group work 2	
L6	25/9	10-12	M33	The Earth's magnetosphere 2, Other magnetospheres	<b>CGF</b> Ch 4.6-4.9, <b>LL</b> Ch V.
L7	30/9	14-16	L51	Aurora, Measurement methods in space plasmas and data analysis 1	<b>CGF</b> Ch 4.5, 10, <b>LL</b> Ch VI, Extra material
T3	3/10	10-12	V22	Mini-group work 3	
L8	7/10	10-12	V22	Space weather and geomagnetic storms	<b>CGF</b> Ch 4.4, <b>LL</b> Ch IV.B-C, VII.A-C
T4	9/10	15-17	Q31	Mini-group work 4	
L9	11/10	10-12	M33	Interstellar and intergalactic plasma, Cosmic radiation, Swedish and international space physics research.	<b>CGF</b> Ch 7-9
T5	15/10	10-12	L51	Mini-group work 5	
L10	16/10	13-15	Q36	Guest lecture: Swedish astronaut Christer Fuglesang	
T6	17/10	15-17	Q31	Round-up	
Written examination	30/10	14-19	B21-24		

# Mini groupwork 1

a)

$$h = \frac{42 \text{ mm}}{7 \text{ mm}} \cdot 6378 \text{ km} \cdot 2 = 77000 \text{ km}$$

The thermal energy is divided into motion in the three dimensions, two of which only give rise to a gyro motion around the magnetic field lines, with the motion along the magnetic field corresponding to an energy

$$E = \frac{k_B T}{2} = \frac{1.38 \cdot 10^{-23} \cdot 1.5 \cdot 10^6}{2} = 1 \cdot 10^{-17} \text{ J}$$

$$v = \sqrt{\frac{2E}{m_e}} = \sqrt{\frac{2 \cdot 10^{-17}}{0.91 \cdot 10^{-30}}} = 4.7 \cdot 10^6 \text{ ms}^{-1}$$

Approximating the loop with a quarter-circle, the electron has to travel a length

$$s = \pi h / 2 = 120\,000 \text{ km}$$

Then we get  $t = 25 \text{ s}$ .

# Energy - temperature

Average energy of molecule/atom:


$$E = \frac{3}{2} k_B T \Rightarrow$$

$$T = \frac{2E}{3k_B}$$

$$1 \text{ eV} = 1.6 \cdot 10^{-19} \text{ J} \Rightarrow$$

$$T = \frac{2E}{3k_B} = \frac{2 \cdot 1.6 \cdot 10^{-19} \text{ J}}{3 \cdot 1.38 \cdot 10^{-23} \frac{\text{J}}{\text{K}}} = 7729 \text{ K}$$

# Gyro motion



## Equipartition principle

Statistically the kinetic energy is equally distributed along the three dimensions:

$$E_{\parallel} = \frac{1}{2} k_B T$$

$$E_{\perp} = \frac{2}{2} k_B T$$

# Mini groupwork 1

b) 
$$f_c = \frac{\omega_c}{2\pi} = \frac{1}{2\pi} \frac{qB}{m} \Rightarrow$$

$$B = \frac{2\pi f_c m}{q} = \frac{2\pi \cdot 1 \cdot 10^{10} \cdot 0.91 \cdot 10^{-30}}{1.6 \cdot 10^{-19}} = 0.36 \text{ T}$$

The perpendicular energy is given by

$$E = 2 \cdot \frac{k_B T}{2} = 2 \cdot \frac{1.38 \cdot 10^{-23} \cdot 1.5 \cdot 10^6}{2} = 2 \cdot 10^{-17} \text{ J}$$


$\Rightarrow$

$$v = \sqrt{\frac{2E}{m_e}} = \sqrt{\frac{4 \cdot 10^{-17}}{0.91 \cdot 10^{-30}}} = 6.6 \cdot 10^6 \text{ ms}^{-1}$$

$$\rho = \frac{m_e v_{\perp}}{qB} = \frac{0.91 \cdot 10^{-30} \cdot 6.6 \cdot 10^6}{1.6 \cdot 10^{-19} \cdot 0.36} = 1.0 \cdot 10^{-4} \text{ m}$$

# Mini groupwork 1

- c) Model the flare by a half torus with minor axis  $r$ , and major axis. From the figure, estimate  $R = 2.6 R_E$ , and  $r = 2 R_E$ .




Let this half-torus be filled with a magnetic field of strength  $B \sim 0.36$  T (using the value in b)). If the volume of the half-torus is  $V$  and the magnetic energy density is  $p_B$ , the total energy is

$$\begin{aligned}
 W &= V p_B = \pi R \pi r^2 \frac{B^2}{2\mu_0} = \pi^2 \cdot 2 \cdot 12^2 R_E^3 \frac{B^2}{2\mu_0} \\
 &= \pi^2 \cdot 2 \cdot 12^2 (6378 \cdot 10^3)^3 \frac{(0.36)^2}{2\mu_0} = 3.8 \cdot 10^{28} \text{ J}
 \end{aligned}$$




# Solar flare mechanism




Electrons are accelerated, collide with solar surface (photosphere) and emit bremsstrahlung (X-rays).

# Solar flare observations



(a) double signature of x-ray emissions at foot of flare



(b) coronal loop filled with hot gas

# Frozen in magnetic flux *PROOF II*

$$\frac{\partial \mathbf{B}}{\partial t} = \underbrace{\nabla \times (\mathbf{v} \times \mathbf{B})}_A + \underbrace{\frac{1}{\mu_0 \sigma} \nabla^2 \mathbf{B}}_B$$

Order of magnitude estimate:

$$\frac{A}{B} = \frac{\nabla \times (\mathbf{v} \times \mathbf{B})}{\frac{1}{\mu_0 \sigma} \nabla^2 \mathbf{B}} \approx \frac{\frac{v \Delta B}{L}}{\frac{\Delta B}{\mu_0 \sigma L^2}} = v L \mu_0 \sigma \equiv R_m$$


Magnetic Reynolds number  $R_m$ :

$$R_m \gg 1 \Rightarrow \frac{\partial \mathbf{B}}{\partial t} = \nabla \times (\mathbf{v} \times \mathbf{B})$$

Frozen-in fields!

$$R_m \ll 1 \Rightarrow \frac{\partial \mathbf{B}}{\partial t} = \frac{1}{\mu_0 \sigma} \nabla^2 \mathbf{B}$$

Diffusion equation!




# Reconnection

In 'diffusion region':

$$R_m = \mu_0 \sigma l v \sim 1$$



Thus: **condition** for frozen-in magnetic field breaks down.

A second **condition** is that there are two regions of magnetic field pointing in *opposite* direction:



- Field lines are “cut” and can be re-connected to other field lines
- **Magnetic energy is transformed into kinetic energy ( $U_o \gg U_i$ )**
- **Plasma from different field lines can mix**

# Reconnection in 1D



$$\frac{\partial \mathbf{B}}{\partial t} = \frac{1}{\mu_0 \sigma} \nabla^2 \mathbf{B} \quad \rightarrow \quad \frac{\partial B_x}{\partial t} = \frac{1}{\mu_0 \sigma} \frac{\partial^2 B_x}{\partial z^2}$$

Diffusion equation! Has solution




$$B_x(z, t) = B_0 \operatorname{erf} \left( \left[ \frac{\mu_0 \sigma}{4t} \right]^{1/2} z \right)$$

The total magnetic energy then decreases with time:

$$W_B = \int_{-\infty}^{\infty} \frac{B^2}{2\mu_0} dz$$

The magnetic energy is converted into heat and kinetic energy in 2D


# Solar flare *energization mechanism*



Two possible reconnection geometries

# Solar wind

Interplanetary current sheet





# Solar wind

## Some basic facts

### Average values

$$n_p = 8 \text{ cm}^{-3}$$


$$v = 320 \text{ km/s}$$

$$T_p = 4 \cdot 10^4 \text{ K}$$

$$T_e = 10^5 \text{ K}$$

$$B = 5 \text{ nT}$$

$$\Phi_K = \rho v^3 / 2 = 0.22 \text{ mW/m}_2$$



# The solar wind today

## Average values

$$n_p = 8 \text{ cm}^{-3}$$

$$v = 320 \text{ km/s}$$


$$T_p = 4 \cdot 10^4 \text{ K}$$

$$T_e = 10^5 \text{ K}$$

$$B = 5 \text{ nT}$$

$$p_D = \rho v^2 / 2 = 0.7 \text{ nPa}$$

$$\Phi_K = \rho v^3 / 2 = 0.22 \text{ mW/m}^2$$



Measurements from ACE spacecraft

<http://www.swpc.noaa.gov/SWN/>

Space Weather Prediction Centre



Guess how long does it take the solar wind to flow from the Sun to the Earth?

Blue

8 min

Yellow

1.5 days

Green

5 hours

Red

5 days



$$t = \frac{s}{v} = \frac{1.496 \cdot 10^{11}}{320 \cdot 10^3} = 467\,500 \text{ s} = 129.9 \text{ h} = 5.4 \text{ days}$$

Red

But maybe

Yellow




if the solar wind is much faster

Does anyone happen to know the mathematical formula for the spiral caused by a rotating garden sprinkler?



# Solar wind


*Magnetic field frozen into solar wind*




This is now seen from "above"! (Looking down on the ecliptic plane from the pole.)

# Solar wind

## *Parker spiral*



# Parker spiral




## *Derivation of $\Psi$ (Parker angle)*

Consider a coordinate system rotating with the sun. The plasma element  $P$  in this coordinate system has two velocity components:  $U_r$  and  $U_\phi$ .

Since the magnetic field is frozen into the solar wind, and follows the orbit of the plasma element  $P$ , at any time  $B$  has to be parallel to  $U$ . Then we have:

$$\tan \psi = \frac{B_\phi}{B_r} = \frac{U_\phi}{U_r} = \left( \frac{\omega r}{u_{SW}} \right)$$



$$d\mathbf{x} = \mathbf{U}_{sw} dt$$




# Solar wind

## Parker spiral

Archimedean spiral:

$$\frac{B_{\phi}}{B_r} = \tan \psi = \left( \frac{\omega r}{u_{SW}} \right)$$



# Archimedean spiral

An Archimedean spiral (also arithmetic spiral), is a spiral named after the 3rd-century-BC Greek mathematician Archimedes; it is the locus of points corresponding to the locations over time of a point moving away from a fixed point with a constant speed along a line which rotates with constant angular velocity. Equivalently, in polar coordinates  $(r, \phi)$  it can be described by the equation (*Wikipedia*)


$$r = a + b\phi$$

$$r = a + b\omega t$$

$$\frac{dr}{dt} = b\omega = u_{SW}$$


$$b = \frac{u_{SW}}{\omega}$$

$$r = R_{sun} + \frac{u_{SW}}{\omega} \phi$$



Use rotation period  
 $T$  of sun:  $T = 27$  days

What is the angle  $\Psi$   
 at Earth's orbit for a  
 typical solar wind  
 speed?



$r = 1 \text{ A.U.}$

Yellow  $\approx 50^\circ$

Red  $\approx 80^\circ$

Blue  $\approx 1^\circ$

Green  $\approx 10^\circ$



$$\Psi = \arctan\left(\frac{\omega r}{u}\right)$$

What is  $\omega$ ?  $\omega = 2\pi f = \frac{2\pi}{T} = \frac{2\pi}{27 \cdot 24 \cdot 60 \cdot 60} = 2.7 \cdot 10^{-6} \text{ s}^{-1}$

$$\Psi = \arctan\left(\frac{\omega r}{u}\right) = \arctan\left(\frac{2.7 \cdot 10^{-6} \cdot 1.5 \cdot 10^{11}}{320 \cdot 10^3}\right) = \arctan(1.27) = 52^\circ$$

Yellow



# Classification of plasmas

- **High density plasmas**

- $\lambda \ll \rho$
- *magnetic field not important, collisions dominate, isotropic.*

- **Medium density plasmas**

- $\rho \ll \lambda \ll l_c$
- *magnetic field important, collisions important, anisotropies.*

- **Low density plasmas**

- $l_c \ll \lambda$
- *magnetic field important, anisotropies, uninhibited motion along magnetic field*

$\rho$ : gyro radius

$\lambda$ : mean free path

$l_c$ : dimension of the plasma



# Plasma models/descriptions

- Single particle motion
- Computer simulations of many-particle dynamics
- Generalization of statistical mechanics (kinetic theory)
- Generalization of fluid mechanics:  
*Magneto-hydrodynamics (MHD)*


# Plasma physics

## *Magnetohydrodynamics (MHD)*

MHD is a combination of


- *fluid-/hydrodynamics* (which is based on Newton's laws of motion)
- *Maxwell's equations* (electrodynamics)

applied on a plasma volume element.



# Magnetohydrodynamics (MHD)

For a volume element of plasma:



$$\mathbf{F} = m\mathbf{a} \quad \Rightarrow$$

$$-\nabla p + n_e q \mathbf{v}_e \times \mathbf{B} + \cancel{\rho q \mathbf{E}} = \rho \frac{d\mathbf{v}}{dt} \quad \Rightarrow$$

quasineutrality

$$(1) \quad \rho \frac{d\mathbf{v}}{dt} = -\nabla p + \mathbf{j} \times \mathbf{B}$$



# Magnetohydrodynamics (MHD)

$$(1) \quad \rho \frac{d\mathbf{v}}{dt} = -\nabla p + \mathbf{j} \times \mathbf{B}$$

This together with two of Maxwell's equations and Ohm's law make up the most common MHD equations:

$$(2) \quad \mathbf{j} = \sigma(\mathbf{E} + \mathbf{v} \times \mathbf{B})$$

$$(3) \quad \nabla \times \mathbf{B} = \mu_0 \left( \mathbf{j} + \epsilon_0 \frac{\partial \mathbf{E}}{\partial t} \right)$$

Only consider slow variations

$$(4) \quad \nabla \times \mathbf{E} = -\frac{\partial \mathbf{B}}{\partial t}$$

# Magnetohydrodynamics (MHD)

$$(1) \quad \rho \frac{d\mathbf{v}}{dt} = -\nabla p + \mathbf{j} \times \mathbf{B}$$

In equilibrium:

$$0 = -\nabla p + \mathbf{j} \times \mathbf{B} \quad \longleftrightarrow$$

$$-\nabla p + \frac{1}{\mu_0} (\nabla \times \mathbf{B}) \times \mathbf{B} = 0$$

$$-\nabla p - \nabla \left( \frac{B^2}{2\mu_0} \right) + \frac{1}{\mu_0} (\mathbf{B} \cdot \nabla) \mathbf{B} = 0$$


Represents tension in magnetic field

If magnetic tension = 0

$$p + \frac{B^2}{2\mu_0} = \textit{konst}$$

Magnetic pressure

# Quasineutrality



$$\Phi = \Phi_0 e^{-x/\lambda_D}$$

Debye length

$$\lambda_D = \sqrt{\frac{\epsilon_0 k_B T}{n_e e^2}}$$


$$\frac{\Delta n}{n} = \frac{(n_e - n_i)}{n_e} < \left( \frac{\lambda_D}{l_c} \right)^2$$

$$l_c \gg \lambda_D \Rightarrow$$

Plasma close to neutral:

$$n_e \approx n_i$$


# Debye lengths






# *The ionosphere*

# Basic principle for creation of ionospheric layer





# Atmospheric scale height

$$-\frac{dp}{dz} = g\rho_m \quad \text{hydrostatic equilibrium for a volume element}$$

$$p = nk_B T = \frac{\rho k_B T}{m} \quad \text{ideal gas law}$$

$$-\frac{k_B T}{m} \frac{d\rho_m}{dz} = g\rho_m \quad \text{if } T \text{ is constant}$$

$$\rho_m = \text{const} \cdot e^{-z/(k_B T/gm)} = \text{const} \cdot e^{-z/H}$$

Scale height

$$H = k_B T/gm$$



## Scale height

$$H = k_B T / gm$$

What is the approximate scale height in the atmosphere right here, right now?

*(0° C = 273 K)*

Blue

1 km

Yellow

30 km

Green

9 km

Red

100 km







$$H = k_B T / gm = (1.38 \cdot 10^{-23} \cdot 290) / (9.81 \cdot 14 \cdot 2 \cdot 1.67 \cdot 10^{-27}) =$$
$$= 8724 \text{ m} \approx 9 \text{ km}$$

Green




What did we neglect  
when we derived the  
scale height?

# Temperature profile



# Atmospheric composition

Longer scale height due to higher temperature



Separate scale heights for different components

Turbulent mixing – one scale height




# Ionosphere

- The ionized, electrically conducting part of the upper atmosphere
- The ionosphere is a **plasma**

# History

- Stewart, 1882: Explained variations in the geomagnetic field
- Kenelly & Heavyside, 1902: explained Marconis transatlantic radio communication experiments
- Appleton & Barnett: experimental proof

# Altitude distribution of electron density ( $n_e$ )



# Continuity equation = conservation of ?

$$\frac{\partial n_e}{\partial t} = q - r - \nabla \cdot (n_e \mathbf{v}_e)$$

Ionization ( $\text{m}^{-3}\text{s}^{-1}$ )

Recombination ( $\text{m}^{-3}\text{s}^{-1}$ )

Flow ( $\text{m}^{-3}\text{s}^{-1}$ )

# Continuity equation

$$\frac{dn_e}{dt} = q - r$$

$$q = a_i I n_n$$

Ionization ( $\text{m}^{-3}\text{s}^{-1}$ )


Recombination ( $\text{m}^{-3}\text{s}^{-1}$ )

$$r = a_r n_e n_i = a_r n_e^2$$

Example:  $e + \text{O}_2^+ \rightarrow \text{O} + \text{O}$  (dissociative recombination)



# UV and X-ray radiation



$$\frac{dI}{dz} = In_n a_a$$



# Derive Chapman layer



# Last Minute!